

The universal R-matrix and factorization of the L-operators related to the Baxter Q-operators

Sergey Khoroshkin* Zengo Tsuboi†

Abstract

We consider the ‘universal monodromy operators’ for the Baxter Q-operators. They are given as images of the universal R-matrix in oscillator representation. We find related universal factorization formulas in $U_q(\hat{sl}(2))$ case.

Keywords: Baxter Q-operator, universal R-matrix, L-operator, integrable system

PACS number: 02.20.Uw, 02.30.Ik

Journal Reference: J. Phys. A: Math. Theor. 47 (2014) 192003 (Fast Track Communications)

doi:10.1088/1751-8113/47/19/192003

1 Introduction

The Baxter Q-operators were introduced by Baxter [1], and are considered to be one of the most powerful tools in quantum integrable systems. The T-operators (transfer matrices) are expressed in terms of Q-operators. Thus Q-operators are fundamental objects.

In particular, Bazhanov, Lukyanov and Zamolodchikov defined Q-operators as the partial trace of the L -operator, which is a bosonic realization of the universal R-matrix, over some q -oscillator representations of the Borel subalgebra of a quantum affine algebra [2]. This ‘ q -oscillator construction’ of the Q-operators was developed and generalized for higher rank algebras, superalgebras and so forth ([3, 4, 5, 6, 7] and references therein). It is known that the q -oscillator representations for the Q-operators are given as limits of some representations of the Borel subalgebra of quantum affine algebra. Recently, a systematic account on this from the point of view of the representation theory was given and used in [8].

The T and Q-operators are given as the traces of (product of) the L-operators, which are images of the universal R-matrix. In particular, L-operators for (infinite dimensional) Verma modules of the quantum affine algebra (or the Yangian) factorize with respect to ‘L-operators for the Q-operators’. Examples for such factorization formulas were given

*Institute of Theoretical and Experimental Physics, 117218 Moscow, Russia; Higher School of Economics, Myasnitskaya 20, 101000, Moscow, Russia

†Present address: Department of Mathematics and Statistics, The University of Melbourne, Royal Parade, Parkville, Victoria 3010, Australia

by a number of authors ([9, 10] and references therein). In this paper, we reconsider this phenomenon in relation to properties of the universal R-matrix and obtain a universal factorization formula, which is independent of the quantum space. Throughout this paper, we use presentations of the universal R-matrix from [12, 13]. We remark that a detailed explanation on the evaluation of the universal R-matrix related to Q-operators, which might be omitted or missing in earlier papers, can be found in a recent series of papers [6].

2 Quantum algebras

2.1 $U_q(\hat{sl}(2))$

The quantum affine algebra $U_q(\hat{sl}(2))$ is a Hopf algebra generated by the generators e_i, f_i, h_i, d , where $i \in \{0, 1\}$. We introduce the q -commutator $[X, Y]_q = XY - qYX$. In particular, $[X, Y]_1 = [X, Y]$. For $i, j \in \{0, 1\}$, the defining relations of the algebra $U_q(\hat{sl}(2))$ are given by

$$\begin{aligned} [h_i, h_j] &= 0, \quad [h_i, e_j] = a_{ij}e_j, \quad [h_i, f_j] = -a_{ij}f_j, \\ [e_i, f_j] &= \delta_{ij} \frac{q^{h_i} - q^{-h_i}}{q - q^{-1}}, \\ [e_i, [e_i, [e_i, e_j]_{q^2}]]_{q^{-2}} &= [f_i, [f_i, [f_i, f_j]_{q^{-2}}]]_{q^2} = 0 \quad i \neq j, \end{aligned} \tag{2.1}$$

where $(a_{ij})_{0 \leq i, j \leq 1}$ is the Cartan matrix

$$(a_{ij})_{0 \leq i, j \leq 1} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}.$$

We use the following co-product $\Delta : U_q(\hat{sl}(2)) \rightarrow U_q(\hat{sl}(2)) \otimes U_q(\hat{sl}(2))$:

$$\begin{aligned} \Delta(e_i) &= e_i \otimes 1 + q^{-h_i} \otimes e_i, \\ \Delta(f_i) &= f_i \otimes q^{h_i} + 1 \otimes f_i, \\ \Delta(h_i) &= h_i \otimes 1 + 1 \otimes h_i. \end{aligned} \tag{2.2}$$

We will also use an opposite co-product defined by

$$\Delta' = \sigma \circ \Delta, \quad \sigma \circ (X \otimes Y) = Y \otimes X, \quad X, Y \in U_q(\hat{sl}(2)). \tag{2.3}$$

We always assume zero value of the central element $h_0 + h_1$. Anti-pode, co-unit and grading element d are not used in this paper.

The Borel subalgebra $\mathcal{B}_+$ (resp. $\mathcal{B}_-$) is generated by e_i, h_i (resp. f_i, h_i), where $i \in \{0, 1\}$. For complex numbers $c_i \in \mathbb{C}$ which obey the relation $\sum_{i=0}^1 c_i = 0$, the transformation

$$h_i \mapsto h_i + c_i, \quad i = 0, 1, \tag{2.4}$$

determines the shift automorphism τ_{c_1} of $\mathcal{B}_+$ (or of $\mathcal{B}_-$). Here we omit the unit element multiplied by the above complex numbers.

There exists a unique element [11, 12] $\mathcal{R}$ in a completion of $\mathcal{B}_+ \otimes \mathcal{B}_-$ called the universal R-matrix which satisfies the following relations

$$\begin{aligned}\Delta'(a) \mathcal{R} &= \mathcal{R} \Delta(a) \quad \text{for } \forall a \in U_q(\hat{sl}(2)), \\ (\Delta \otimes 1) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{23}, \\ (1 \otimes \Delta) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{12}\end{aligned}\tag{2.5}$$

where $\mathcal{R}_{12} = \mathcal{R} \otimes 1$, $\mathcal{R}_{23} = 1 \otimes \mathcal{R}$, $\mathcal{R}_{13} = (\sigma \otimes 1) \mathcal{R}_{23}$. The Yang-Baxter equation

$$\mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23} = \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12}, \tag{2.6}$$

is a corollary of these relations (2.5). The universal R-matrix can be written in the form

$$\mathcal{R} = \overline{\mathcal{R}} q^{\mathcal{K}}, \quad \mathcal{K} = \frac{1}{2} h_1 \otimes h_1. \tag{2.7}$$

Here $\overline{\mathcal{R}}$ is the reduced universal R-matrix, which is a series in $e_j \otimes 1$ and $1 \otimes f_j$ and does not contain Cartan elements (to denote this symbolically, we will use a notation $\overline{\mathcal{R}} = \overline{\mathcal{R}}(\{e_0 \otimes 1, e_1 \otimes 1, 1 \otimes f_0, 1 \otimes f_1\})$). Thus the reduced universal R-matrix is unchanged under the shift automorphism τ_{c_1} of $\mathcal{B}_+$, see (2.4), while the prefactor $\mathcal{K}$ is shifted as

$$\mathcal{K} \mapsto \mathcal{K} + \frac{c_1}{2} (1 \otimes h_1). \tag{2.8}$$

2.2 $U_q(sl(2))$

The algebra $U_q(sl(2))$ is generated by the elements E, F, H . The defining relations are

$$\begin{aligned}[H, E] &= 2E, & [H, F] &= -2F, \\ [E, F] &= \frac{q^H - q^{-H}}{q - q^{-1}}.\end{aligned}\tag{2.9}$$

The following elements are central in $U_q(sl(2))$:

$$C^{(r)} = FE + \frac{q^{H+1} + q^{-H-1}}{(q - q^{-1})^2}, \quad C^{(l)} = EF + \frac{q^{H-1} + q^{-H+1}}{(q - q^{-1})^2} \tag{2.10}$$

Due to the relations (2.9), $C^{(r)} = C^{(l)}$ holds.

There is an evaluation map $\text{ev}_x: U_q(\hat{sl}(2)) \mapsto U_q(sl(2))$:

$$\begin{aligned}e_0 &\mapsto F, & f_0 &\mapsto E, & h_0 &\mapsto -H, \\ e_1 &\mapsto xE, & f_1 &\mapsto x^{-1}F, & h_1 &\mapsto H,\end{aligned}\tag{2.11}$$

where $x \in \mathbb{C}^*$ is a parameter. Let π_μ^+ be the Verma module over $U_q(sl(2))$ with the highest weight μ . In a basis $\{v_n | n \in \mathbb{Z}_{\geq 0}\}$, we have

$$Hv_n = (\mu - 2n)v_n, \quad Ev_n = [n]_q [\mu - n + 1]_q v_{n-1}, \quad Fv_n = v_{n+1}. \tag{2.12}$$

For $\mu \in \mathbb{Z}_{\geq 0}$, the module π_μ^+ contains an invariant subspace isomorphic to $\pi_{-\mu-2}^+$, and the quotient space $\pi_\mu^+/\pi_{-\mu-2}^+$ is isomorphic to the finite dimensional irreducible module π_μ with the highest weight μ . In particular, $\pi_1(E) = E_{12}$, $\pi_1(F) = E_{21}$ and $\pi_1(H) = E_{11} - E_{22}$ gives the fundamental representation of $U_q(sl(2))$, where E_{ij} is a 2×2 matrix unit whose (k, l) -element is $\delta_{i,k}\delta_{j,l}$.

Then the compositions $\pi_\mu^+(x) = \pi_\mu^+ \circ \mathbf{ev}_x$ and $\pi_\mu(x) = \pi_\mu \circ \mathbf{ev}_x$ give evaluation representations of $U_q(\hat{sl}(2))$.

2.3 q-oscillator algebras

We introduce two kinds of oscillator algebras Osc_i ($i = 1, 2$). They are generated by the elements $\mathbf{h}_i, \mathbf{e}_i, \mathbf{f}_i$ which are subject to the relations:

$$\begin{aligned} [\mathbf{h}_1, \mathbf{h}_1] &= 0, & [\mathbf{h}_1, \mathbf{e}_1] &= 2\mathbf{e}_1, & [\mathbf{h}_1, \mathbf{f}_1] &= -2\mathbf{f}_1, \\ \mathbf{f}_1\mathbf{e}_1 &= q \frac{1 - q^{\mathbf{h}_1}}{(q - q^{-1})^2}, & \mathbf{e}_1\mathbf{f}_1 &= q \frac{1 - q^{\mathbf{h}_1-2}}{(q - q^{-1})^2}, \end{aligned} \quad (2.13)$$

$$\begin{aligned} [\mathbf{h}_2, \mathbf{h}_2] &= 0, & [\mathbf{h}_2, \mathbf{e}_2] &= 2\mathbf{e}_2, & [\mathbf{h}_2, \mathbf{f}_2] &= -2\mathbf{f}_2, \\ \mathbf{f}_2\mathbf{e}_2 &= q^{-1} \frac{1 - q^{-\mathbf{h}_2}}{(q - q^{-1})^2}, & \mathbf{e}_2\mathbf{f}_2 &= q^{-1} \frac{1 - q^{-\mathbf{h}_2+2}}{(q - q^{-1})^2}, \end{aligned} \quad (2.14)$$

Note that Osc_1 and Osc_2 can be swapped by the transformation $q \mapsto q^{-1}$. One can derive the following corollaries of (2.13) and (2.14):

$$[\mathbf{e}_1, \mathbf{f}_1] = \frac{q^{\mathbf{h}_1}}{q - q^{-1}}, \quad [\mathbf{e}_2, \mathbf{f}_2] = -\frac{q^{-\mathbf{h}_2}}{q - q^{-1}}, \quad (2.15)$$

$$[\mathbf{e}_1, \mathbf{f}_1]_{q^{-2}} = \frac{1}{q - q^{-1}}, \quad [\mathbf{e}_2, \mathbf{f}_2]_{q^2} = -\frac{1}{q - q^{-1}}. \quad (2.16)$$

Relations (2.15) are nothing but contractions¹ of the relations (2.9). On the other hand, the relations (2.16) are conditions that central elements take constant values. The following limits of the generators of $U_q(sl(2))$ for the Verma module π_μ^+

$$\begin{aligned} \lim_{q^{-\mu} \rightarrow 0} (H - \mu) &= \mathbf{h}_1, & \lim_{q^{-\mu} \rightarrow 0} q^{-H-\mu} &= 0, \\ \lim_{q^{-\mu} \rightarrow 0} F &= \mathbf{f}_1, & \lim_{q^{-\mu} \rightarrow 0} Eq^{-\mu} &= \mathbf{e}_1, \end{aligned} \quad (2.17)$$

$$\begin{aligned} \lim_{q^\mu \rightarrow 0} (H - \mu) &= \mathbf{h}_2, & \lim_{q^\mu \rightarrow 0} q^{H+\mu} &= 0, \\ \lim_{q^\mu \rightarrow 0} F &= \mathbf{f}_2, & \lim_{q^\mu \rightarrow 0} Eq^\mu &= \mathbf{e}_2, \end{aligned} \quad (2.18)$$

realize the q-oscillator algebras Osc_1 and Osc_2 , respectively. By this construction, oscillator algebras Osc_1 and Osc_2 are given together with irreducible highest weight representations

¹Contractions of a quantum algebra and its relation to a q-oscillator algebra was discussed in [14].

W_1 and W_2 of zero highest weight. The vector space of W_1 is generated by vectors $v_n^{(1)} = \mathbf{f}_1^n v_0^{(1)}$, and the vector space of W_2 is generated by vectors $v_n^{(2)} = \mathbf{f}_2^n v_0^{(2)}$, so that

$$\mathbf{e}_1 v_n^{(1)} = [n]_q \frac{q^{-n+1}}{q - q^{-1}} v_{n-1}^{(1)}, \quad \mathbf{e}_2 v_n^{(2)} = -[n]_q \frac{q^{n-1}}{q - q^{-1}} v_{n-1}^{(2)}.$$

Here $[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}$.

3 L-operators

In the following we denote $\lambda = q - q^{-1}$. We set

$$\mathbf{L}(x) = \begin{pmatrix} q^{\frac{H}{2}} - q^{-1} x q^{-\frac{H}{2}} & \lambda F q^{-\frac{H}{2}} \\ \lambda x E q^{\frac{H}{2}} & q^{-\frac{H}{2}} - q^{-1} x q^{\frac{H}{2}} \end{pmatrix}, \quad (3.1)$$

We have

$$q \det \mathbf{L}(x) = \mathbf{L}_{11}(x) \mathbf{L}_{22}(q^{-2}x) - q^{-1} \mathbf{L}_{21}(x) \mathbf{L}_{12}(q^{-2}x) = 1 - q^{-2} \lambda^2 C^{(l)} x + q^{-4} x^2. \quad (3.2)$$

Then

$$\mathbf{L}(x) = \phi(x)(\mathbf{e}v_x \otimes \pi_1(1))\mathcal{R},$$

and $\phi(x)$ solves the relation (3.2), so that $\phi(x)\phi(q^{-2}x) = 1 - q^{-2} \lambda^2 C^{(l)} x + q^{-4} x^2$. Explicitly, it implies $\phi(x) = \exp(\sum_{k=1}^{\infty} \frac{-C_k}{1+q^{2k}} \frac{x^k}{k})$, with central elements $C_k \in U_q(\hat{sl}(2))$, so that $\pi_{\mu}^{+}(C_k) = q^{(\mu+1)k} + q^{-(\mu+1)k}$.

Define $\mathbf{L}^{(1)}(x)$ and $\mathbf{L}^{(2)}(x)$ as the following limits of homomorphic images of $\mathbf{L}(x)$:

$$\mathbf{L}^{(1)}(x) = \lim_{q^{-\mu} \rightarrow 0} (\pi_{\mu}^{+} \otimes 1) \mathbf{L}(x q^{-\mu}) q^{-\frac{\mu \otimes \pi_1(1)(h_1)}{2}} = \begin{pmatrix} q^{\frac{h_1}{2}} & \lambda \mathbf{f}_1 q^{-\frac{h_1}{2}} \\ \lambda x \mathbf{e}_1 q^{\frac{h_1}{2}} & q^{-\frac{h_1}{2}} - q^{-1} x q^{\frac{h_1}{2}} \end{pmatrix}, \quad (3.3)$$

$$\mathbf{L}^{(2)}(x) = \lim_{q^{\mu} \rightarrow 0} (\pi_{\mu}^{+} \otimes 1) \mathbf{L}(x q^{\mu}) q^{-\frac{\mu \otimes \pi_1(1)(h_1)}{2}} = \begin{pmatrix} q^{\frac{h_2}{2}} - q^{-1} x q^{-\frac{h_2}{2}} & \lambda \mathbf{f}_2 q^{-\frac{h_2}{2}} \\ \lambda x \mathbf{e}_2 q^{\frac{h_2}{2}} & q^{-\frac{h_2}{2}} \end{pmatrix}, \quad (3.4)$$

where the limits (2.17) and (2.18) are applied² correspondingly to (3.1). The latter L -operators can be also presented as homomorphic images of the universal R -matrix under the homomorphisms $\rho_x^{(i)} : \mathcal{B}_+ \rightarrow \text{Osc}_i$ defined by the relations

$$\rho_x^{(i)}(e_0) = \mathbf{f}_i, \quad \rho_x^{(i)}(e_1) = x \mathbf{e}_i, \quad \rho_x^{(i)}(h_0) = -\mathbf{h}_i, \quad \rho_x^{(i)}(h_1) = \mathbf{h}_i, \quad i = 1, 2. \quad (3.5)$$

Namely we have the relations $\mathbf{L}^{(i)}(x) = \phi^{(i)}(x)(\rho_x^{(i)} \otimes \pi_1(1))\mathcal{R}$, where

$$\phi^{(1)}(x) = \exp\left(\sum_{k=1}^{\infty} \frac{-q^k}{1+q^{2k}} \frac{x^k}{k}\right), \quad \phi^{(2)}(x) = \exp\left(\sum_{k=1}^{\infty} \frac{-q^{-k}}{1+q^{2k}} \frac{x^k}{k}\right).$$

²The factor $q^{-\frac{\mu \otimes \pi_1(1)(h_1)}{2}}$ came from (2.8) for $c_1 = -\mu$.

The L-operators (3.3) and (3.4) are given by some contraction³ of the original L-operator (3.1) with respect to the Cartan part (diagonal elements). We will also use notations⁴

$$\mathcal{L}^{(\mu,+)}(x) = (\pi_\mu^+(x) \otimes 1)\mathcal{R} \quad \text{and} \quad \mathcal{A}^{(i)}(x) = (\rho_x^{(i)} \otimes 1)\mathcal{R}.$$

Thus $\mathbf{L}^{(i)}(x) = \phi^{(i)}(x)(1 \otimes \pi_1(1))\mathcal{A}^{(i)}(x)$.

4 Representations of $\mathcal{B}_+$ and the factorization formula

In this section, we reconsider the q-oscillator representations of $\mathcal{B}_+$ introduced in [2] (see also a detailed explanation in [4]), and give a factorization formula of the L-operator. We will use the co-product in [13, 12] rather than the one in [2]. We will repeatedly use formulas in Appendix B.

Denote by $W_i(x)$ the pullbacks of basic highest weight representations of the oscillator algebras by means of the homomorphisms $\rho_x^{(i)} : \mathcal{B}_+ \rightarrow \text{Osc}_i$. These are representations of $\mathcal{B}_+$ in the spaces W_i . In tensor product $W_1(xq^\mu) \otimes W_2(xq^{-\mu})$ we consider the basis $v_{n,m}$

$$v_{n,m} = \exp_{q^{-2}}((q - q^{-1})\mathbf{e}_1 \otimes \mathbf{f}_2) v_n^{(1)} \otimes v_m^{(2)} = \sum_{k=0}^n \binom{n}{k}_{q^{-2}} v_{n-k}^{(1)} \otimes v_{m+k}^{(2)}, \quad (4.1)$$

where we introduce notations

$$\begin{aligned} \exp_q(x) &= 1 + \sum_{k=1}^{\infty} \frac{x^k}{(k)_q!}, \quad \binom{n}{k}_q = \frac{(n)_q!}{(k)_q!(n-k)_q!}, \\ (k)_q! &= (1)_q(2)_q \cdots (k)_q, \quad (k)_q = \frac{1 - q^k}{1 - q}. \end{aligned}$$

The action of the generators of $\mathcal{B}_+$ on these vectors is as follows:

$$\begin{aligned} \Delta(e_0)v_{n,m} &= v_{n+1,m}, \\ \Delta(e_1)v_{n,m} &= x[n]_q[\mu - n + 1]_q v_{n-1,m} - \frac{xq^{-\mu+2n+m-1}[m]_q}{q - q^{-1}} v_{n,m-1}, \\ \Delta(h_0)v_{n,m} &= 2(n+m)v_{n,m}, \quad \Delta(h_1)v_{n,m} = -2(n+m)v_{n,m}. \end{aligned} \quad (4.2)$$

Here we see a filtration by m , with quotients isomorphic to evaluation Verma modules $\pi_\mu^+(x)$ shifted by automorphisms τ_{-2m} . Set $E_1 = (q^\mu - q^{-\mu-\mathbf{h}_1})\mathbf{e}_1$, $F_1 = \mathbf{f}_1$, $H_1 = \mathbf{h}_1 + \mu$, which realize the Verma module π_μ^+ on the basis $\{v_n^{(1)} | n \in \mathbb{Z}_{\geq 0}\}$. In particular, E_1, F_1 and H_1 satisfy $U_q(sl(2))$ relations (2.9). We also set $E_2 = \mathbf{e}_2$, $H_2 = \mathbf{h}_2 - \mu$.

³A preliminary form of such contraction was previously discussed in [15] and developed in [16, 7].

⁴We find that $\mathcal{A}^{(2)}(x)$ contains an infinite product of q-exponentials, while $\mathcal{A}^{(1)}(x)$ contains only finite (two) for the non-Cartan part.

Then, the above relations (4.1) on vectors $v_n^{(1)} \otimes v_m^{(2)}$ can be rewritten as the following identities on operators in $W_1 \otimes W_2$

$$\begin{aligned} \mathbf{O}^{-1} \Delta(e_0) \mathbf{O} &= F_1 \otimes 1, & \mathbf{O}^{-1} \Delta(e_1) \mathbf{O} &= xE_1 \otimes 1 + q^{-H_1} \otimes xE_2, \\ \mathbf{O}^{-1} \Delta(h_0) \mathbf{O} &= -H_1 \otimes 1 - 1 \otimes H_2, & \mathbf{O}^{-1} \Delta(h_1) \mathbf{O} &= H_1 \otimes 1 + 1 \otimes H_2, \end{aligned} \quad (4.3)$$

where we set $\mathbf{O} = \exp_{q^{-2}}(\lambda \mathbf{e}_1 \otimes \mathbf{f}_2)$. This should be interpreted as

$$\mathbf{O}^{-1} \left((\rho_{xq^\mu}^{(1)} \otimes \rho_{xq^{-\mu}}^{(2)}) \Delta(\mathbf{a}) \right) \mathbf{O} = (\text{ev}_x^{(1)} \otimes \text{ev}_x^{(2)}) \Delta(\mathbf{a}), \quad (4.4)$$

for $\mathbf{a} \in \mathcal{B}_+$, where we introduced the evaluation maps of $\mathcal{B}_+$ into two algebras, which may be regarded as certain quotients of $U_q(sl(2))$:

$$\text{ev}_x^{(1)}(e_0) = F_1, \quad \text{ev}_x^{(1)}(e_1) = xE_1, \quad \text{ev}_x^{(1)}(h_0) = -H_1, \quad \text{ev}_x^{(1)}(h_1) = H_1, \quad (4.5)$$

$$\text{ev}_x^{(2)}(e_0) = 0, \quad \text{ev}_x^{(2)}(e_1) = xE_2, \quad \text{ev}_x^{(2)}(h_0) = -H_2, \quad \text{ev}_x^{(2)}(h_1) = H_2. \quad (4.6)$$

Let us evaluate the universal R-matrix $(\Delta \otimes 1)\mathcal{R} = \mathcal{R}_{13}\mathcal{R}_{23}$ on $W_1(xq^\mu) \otimes W_2(xq^{-\mu}) \otimes 1$.

$$\begin{aligned} \mathbf{O}_{12}^{-1} \left((\rho_{xq^\mu}^{(1)} \otimes \rho_{xq^{-\mu}}^{(2)} \otimes 1)(\Delta \otimes 1)\mathcal{R} \right) \mathbf{O}_{12} &= \mathbf{O}_{12}^{-1} \left((\rho_{xq^\mu}^{(1)} \otimes \rho_{xq^{-\mu}}^{(2)} \otimes 1)\mathcal{R}_{13}\mathcal{R}_{23} \right) \mathbf{O}_{12} \\ &= \mathbf{O}_{12}^{-1} \mathcal{A}_{13}^{(1)}(xq^\mu) \mathcal{A}_{23}^{(2)}(xq^{-\mu}) \mathbf{O}_{12}. \end{aligned} \quad (4.7)$$

On the other hand, taking note on (4.4), we obtain

$$\begin{aligned} \mathbf{O}_{12}^{-1} \left((\rho_{xq^\mu}^{(1)} \otimes \rho_{xq^{-\mu}}^{(2)} \otimes 1)(\Delta \otimes 1)\mathcal{R} \right) \mathbf{O}_{12} &= (\text{ev}_x^{(1)} \otimes \text{ev}_x^{(2)} \otimes 1)(\Delta \otimes 1)\mathcal{R} \\ &= (\text{ev}_x^{(1)} \otimes \text{ev}_x^{(2)} \otimes 1)\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{L}_{13}^{(\mu,+)}(x) \left((1 \otimes \text{ev}_x^{(2)} \otimes 1)\mathcal{R}_{23} \right), \end{aligned} \quad (4.8)$$

Combining (4.7) and (4.8), we obtain the factorization formula

$$\mathbf{O}_{12}^{-1} \mathcal{A}_{13}^{(1)}(xq^\mu) \mathcal{A}_{23}^{(2)}(xq^{-\mu}) \mathbf{O}_{12} = \mathcal{L}_{13}^{(\mu,+)}(x) \left((1 \otimes \text{ev}_x^{(2)} \otimes 1)\mathcal{R}_{23} \right). \quad (4.9)$$

Note that the co-product of the universal R-matrix has the form $(\Delta \otimes 1)\mathcal{R} = \overline{\mathcal{R}}(\{\Delta(e_0) \otimes 1, \Delta(e_1) \otimes 1, 1 \otimes 1 \otimes f_0, 1 \otimes 1 \otimes f_1\})q^{\frac{\Delta(h_1) \otimes h_1}{2}}$. Then we obtain

$$\begin{aligned} (1 \otimes \text{ev}_x^{(2)} \otimes 1)\mathcal{R}_{23} &= \overline{\mathcal{R}}(\{0, 1 \otimes x\mathbf{e}_2 \otimes 1, 1 \otimes 1 \otimes f_0, 1 \otimes 1 \otimes f_1\})q^{\frac{1 \otimes (\mathbf{h}_2 - \mu) \otimes h_1}{2}} = \\ &= \exp_{q^{-2}}(\lambda \otimes x\mathbf{e}_2 \otimes f_1) q^{\frac{1 \otimes (\mathbf{h}_2 - \mu) \otimes h_1}{2}}, \end{aligned} \quad (4.10)$$

where we used an explicit form of the universal R-matrix [12, 13]. The universal R-matrix is an infinite product of q-exponentials. However, only one of them is non-trivial in this case since the image of the generator e_0 is 0. From (4.9), we arrive at the universal factorization formula in $\mathcal{B}_+$, which is independent on the quantum space (which is a representation of $\mathcal{B}_-$).

Theorem 4.1.

$$\mathbf{O}_{12}^{-1} \mathcal{A}_{13}^{(1)}(xq^\mu) \mathcal{A}_{23}^{(2)}(xq^{-\mu}) \mathbf{O}_{12} = \mathcal{L}_{13}^{(\mu,+)}(x) \exp_{q^{-2}}(\lambda \otimes x\mathbf{e}_2 \otimes f_1) q^{\frac{1 \otimes (\mathbf{h}_2 - \mu) \otimes h_1}{2}}. \quad (4.11)$$

In the right hand side, the first factor says that we have an evaluation Verma module with highest weight μ , the second factor says that we have a filtration indexed by the powers of $\mathbf{e}_2$ and the third factor says that in each quotient of the filtration the evaluation Verma module is given with a shift automorphism $\tau_{-\mu}$ of $\mathcal{B}_+$. A similar factorization will also occur for $\mathcal{B}_-$ (see Appendix A). Let us evaluate the quantum space (the third space) of (4.11) to the fundamental evaluation representation $\pi_1(1)$ of $\mathcal{B}_-$ ($f_0 = E_{12}, f_1 = E_{21}, h_1 = -h_0 = E_{11} - E_{22}$). Then we obtain the following factorization formula for the L-operators (3.1) for π_μ^+ , (3.3) and (3.4).

Corollary 4.2.

$$\begin{aligned} \mathbf{O}_{12}^{-1} \mathbf{L}_{13}^{(1)}(xq^\mu) \mathbf{L}_{23}^{(2)}(xq^{-\mu}) \mathbf{O}_{12} = \\ = \begin{pmatrix} q^{\frac{H_1}{2}} - q^{-1}xq^{-\frac{H_1}{2}} & \lambda F_1 q^{-\frac{H_1}{2}} \\ \lambda x E_1 q^{\frac{H_1}{2}} & q^{-\frac{H_1}{2}} - q^{-1}xq^{\frac{H_1}{2}} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \lambda x \mathbf{e}_2 & 1 \end{pmatrix} \begin{pmatrix} q^{\frac{h_2 - \mu}{2}} & 0 \\ 0 & q^{-\frac{h_2 - \mu}{2}} \end{pmatrix}. \end{aligned} \quad (4.12)$$

Here we use the identity

$$\pi_\mu^+ \phi(x) = \phi^{(1)}(q^\mu x) \phi^{(2)}(q^{-\mu} x).$$

A factorization formula for the rational case ($q = 1$), which is similar to (4.12), was given in [10].

5 Baxter Q-operators

In this section, we define T and Q-operators, and mention a factorization formula for the T-operators. We will use a convention of [7] (or [5]). We define the universal boundary transformation as

$$\mathcal{D} = q^{\varphi h_1}, \quad \varphi \in \mathbb{C}. \quad (5.1)$$

The commutativity $(\mathcal{D} \otimes \mathcal{D})\mathcal{R} = \mathcal{R}(\mathcal{D} \otimes \mathcal{D})$ follows from (2.2). It implies the YB relation for the modified R -matrix $\mathcal{R}(\mathcal{D} \otimes 1)$.

We define the universal T-operator for the Verma module as

$$\mathcal{T}_\mu^+(x) = (\text{Tr}_{\pi_\mu^+(x)} \otimes 1) [\mathcal{R}(\mathcal{D} \otimes 1)] = (\text{Tr}_{\pi^+(x)} \otimes 1) [\overline{\mathcal{R}} \overline{\mathcal{D}}], \quad (5.2)$$

where

$$\overline{\mathcal{D}} = q^{\mathcal{K}}(\mathcal{D} \otimes 1) = (1 \otimes z)^{h_1 \otimes 1}, \quad z = q^{\frac{1}{2}h_1 + \varphi}. \quad (5.3)$$

We also define the universal Q-operators as

$$\mathcal{Q}^{(i)}(x) = \chi_i^{-1}(\text{Tr}_{W_i(x)} \otimes 1) [\mathcal{R}(\mathcal{D} \otimes 1)] = \chi_i^{-1}(\text{Tr}_{W_i(x)} \otimes 1) [\overline{\mathcal{R}} \overline{\mathcal{D}}] \quad i = 1, 2, \quad (5.4)$$

where we introduce normalization operators

$$\chi_i = (\text{Tr}_{W_i(x)} \otimes 1) [\overline{\mathcal{D}}] = \frac{1}{1 - z^{-2}}. \quad (5.5)$$

Eq. (5.4) also have an expression $\mathcal{Q}^{(i)}(x) = \chi_i^{-1}(\text{Tr}_{W_i} \otimes 1) \left[\mathcal{A}^{(i)}(x) \rho_x^{(i)}(\mathcal{D}) \otimes 1 \right]$. Let us apply (4.4) for $\mathcal{D}$, multiply this by (4.11) and take the trace over the first and the second space. Then taking note on the definitions (5.2) and (5.4), we arrive at the factorization formula on the T-operator.

Theorem 5.1. (cf. [2])

$$\mathcal{T}_\mu^+(x) = \frac{z^\mu}{1 - z^{-2}} \mathcal{Q}^{(1)}(xq^\mu) \mathcal{Q}^{(2)}(xq^{-\mu}) \quad (5.6)$$

Note that $\mathcal{T}_\mu^+(0) = z^\mu/(1 - z^{-2})$ is the character of the Verma module π_μ^+ . Thus the T-operator is the Baxterization of the character by the Baxter Q-operators in the sense of [17, 7].

Appendix A: various factorization formulas

Representations of $\mathcal{B}_-$

Set $F_2 = \mathbf{f}_2(q^\mu - q^{-\mu+\mathbf{h}_2})$, $E_2 = \mathbf{e}_2$, $H_2 = \mathbf{h}_2 - \mu$. Let us introduce evaluation maps from $\mathcal{B}_-$

$$\text{ev}_x^{(1)}(f_0) = 0, \quad \text{ev}_x^{(1)}(f_1) = x^{-1}F_1, \quad \text{ev}_x^{(1)}(h_0) = -H_1, \quad \text{ev}_x^{(1)}(h_1) = H_1, \quad (A1)$$

$$\text{ev}_x^{(2)}(f_0) = E_2, \quad \text{ev}_x^{(2)}(f_1) = x^{-1}F_2, \quad \text{ev}_x^{(2)}(h_0) = -H_2, \quad \text{ev}_x^{(2)}(h_1) = H_2, \quad (A2)$$

$$\rho_x^{(i)}(f_0) = \mathbf{e}_i, \quad \rho_x^{(i)}(f_1) = x^{-1}\mathbf{f}_i, \quad \rho_x^{(i)}(h_0) = -\mathbf{h}_i, \quad \rho_x^{(i)}(h_1) = \mathbf{h}_i, \quad i = 1, 2. \quad (A3)$$

Then we find that (4.4) also⁵ hold true for any $\mathbf{a} \in \mathcal{B}_-$ under (A1)-(A3). From a similar argument on $\mathcal{B}_+$ case, we arrive at the universal factorization formula on $\mathcal{B}_-$.

$$\begin{aligned} \mathbf{O}_{23}^{-1} \left((1 \otimes 1 \otimes \rho_{xq^{-\mu}}^{(2)}) \mathcal{R}_{13} \right) \left((1 \otimes \rho_{xq^\mu}^{(1)} \otimes 1) \mathcal{R}_{12} \right) \mathbf{O}_{23} = \\ = ((1 \otimes 1 \otimes \text{ev}_x^{(2)}) \mathcal{R}_{13}) \exp_{q^{-2}} (\lambda e_1 \otimes x^{-1}\mathbf{f}_1 \otimes 1) q^{\frac{h_1 \otimes (\mathbf{h}_1 + \mu) \otimes 1}{2}}. \end{aligned} \quad (A4)$$

The opposite case $W_2(xq^{-\mu}) \otimes W_1(xq^\mu)$

Set

$$\tilde{E}_1 = (q^{-\mu} - q^{\mu-\mathbf{h}_1})\mathbf{e}_1, \quad \tilde{F}_1 = \mathbf{f}_1, \quad \tilde{H}_1 = \mathbf{h}_1 - \mu, \quad (A5)$$

$$\tilde{E}_2 = \mathbf{e}_2, \quad \tilde{F}_2 = \mathbf{f}_2(q^{-\mu} - q^{\mu+\mathbf{h}_2}), \quad \tilde{H}_2 = \mathbf{h}_2 + \mu. \quad (A6)$$

These generators with tilde can be obtained from the ones without tilde by $\mu \rightarrow -\mu$. Let us introduce extended evaluation maps

$$\tilde{\text{ev}}_x^{(1)}(e_0) = \tilde{F}_1, \quad \tilde{\text{ev}}_x^{(1)}(e_1) = 0, \quad \tilde{\text{ev}}_x^{(1)}(h_0) = -\tilde{H}_1, \quad \tilde{\text{ev}}_x^{(1)}(h_1) = \tilde{H}_1, \quad (A7)$$

$$\tilde{\text{ev}}_x^{(1)}(f_0) = \tilde{E}_1, \quad \tilde{\text{ev}}_x^{(1)}(f_1) = x^{-1}\tilde{F}_1, \quad (A8)$$

$$\tilde{\text{ev}}_x^{(2)}(e_0) = \tilde{F}_2, \quad \tilde{\text{ev}}_x^{(2)}(e_1) = x\tilde{E}_2, \quad \tilde{\text{ev}}_x^{(2)}(h_0) = -\tilde{H}_2, \quad \tilde{\text{ev}}_x^{(2)}(h_1) = \tilde{H}_2, \quad (A9)$$

$$\tilde{\text{ev}}_x^{(2)}(f_0) = \tilde{E}_2, \quad \tilde{\text{ev}}_x^{(2)}(f_1) = 0. \quad (A10)$$

⁵However, this does not mean that the maps (A3) can be extended to the whole algebra $U_q(\hat{sl}(2))$. They are regarded as maps from contracted algebras on $U_q(\hat{sl}(2))$ (cf. [7]).

Then we obtain

$$\mathbf{O}_{21}^{-1} q^{-\frac{\mu(1 \otimes \mathbf{h}_1 - \mathbf{h}_2 \otimes 1)}{2}} \left((\rho_{xq^{-\mu}}^{(2)} \otimes \rho_{xq^\mu}^{(1)}) \Delta(\mathbf{a}) \right) q^{\frac{\mu(1 \otimes \mathbf{h}_1 - \mathbf{h}_2 \otimes 1)}{2}} \mathbf{O}_{21} = (\tilde{\mathbf{e}}\mathbf{v}_x^{(2)} \otimes \tilde{\mathbf{e}}\mathbf{v}_x^{(1)}) \Delta(\mathbf{a}),$$

$$\mathbf{a} \in \mathcal{B}_+ \quad \text{or} \quad \mathbf{a} \in \mathcal{B}_-. \quad (\text{A11})$$

Let us evaluate the universal R-matrix $(\Delta \otimes 1)\mathcal{R} = \mathcal{R}_{13}\mathcal{R}_{23}$ on $W_2(xq^{-\mu}) \otimes W_1(xq^\mu) \otimes 1$. From (A11), we arrive at the universal factorization formula on $\mathcal{B}_+$

$$\mathbf{O}_{21}^{-1} q^{-\frac{\mu(1 \otimes \mathbf{h}_1 - \mathbf{h}_2 \otimes 1) \otimes 1}{2}} \left((\rho_{xq^{-\mu}}^{(2)} \otimes 1 \otimes 1) \mathcal{R}_{13} \right) \cdot \left((1 \otimes \rho_{xq^\mu}^{(1)} \otimes 1) \mathcal{R}_{23} \right) q^{\frac{\mu(1 \otimes \mathbf{h}_1 - \mathbf{h}_2 \otimes 1) \otimes 1}{2}} \mathbf{O}_{21} =$$

$$= ((\tilde{\mathbf{e}}\mathbf{v}_x^{(2)} \otimes 1 \otimes 1) \mathcal{R}_{13}) \exp_{q^{-2}}(\lambda \otimes \mathbf{f}_1 \otimes f_0) q^{\frac{1 \otimes (\mathbf{h}_1 - \mu) \otimes h_1}{2}}. \quad (\text{A12})$$

Factorization on both $\mathcal{B}_+$ and $\mathcal{B}_-$

We expect the following factorization formula with respect to four kinds of images of the universal R-matrix:

$$\mathbf{O}_{34}^{-1} \mathbf{O}_{12}^{-1} \left((\rho_{xq^\mu}^{(1)} \otimes 1 \otimes 1 \otimes \rho_{yq^{-\nu}}^{(2)}) \mathcal{R}_{14} \right) \left((\rho_{xq^\mu}^{(1)} \otimes 1 \otimes \rho_{yq^\nu}^{(1)} \otimes 1) \mathcal{R}_{13} \right) \times$$

$$\times \left((1 \otimes \rho_{xq^{-\mu}}^{(2)} \otimes 1 \otimes \rho_{yq^{-\nu}}^{(2)}) \mathcal{R}_{24} \right) \left((1 \otimes \rho_{xq^{-\mu}}^{(2)} \otimes \rho_{yq^\nu}^{(1)} \otimes 1) \mathcal{R}_{23} \right) \mathbf{O}_{12} \mathbf{O}_{34} =$$

$$= ((\mathbf{e}\mathbf{v}_x^{(1)} \otimes 1 \otimes 1 \otimes \mathbf{e}\mathbf{v}_y^{(2)}) \mathcal{R}_{14}) \exp_{q^{-2}}(\lambda x(q^\mu - q^{-\mu - \mathbf{h}_1}) \mathbf{e}_1 \otimes 1 \otimes y^{-1} \mathbf{f}_1 \otimes 1)$$

$$\times \exp_{q^{-2}}(\lambda \otimes x \mathbf{e}_2 \otimes 1 \otimes y^{-1} \mathbf{f}_2(q^\nu - q^{-\nu + \mathbf{h}_2})) \exp_{q^{-2}}(\lambda q^{-\mathbf{h}_1 - \mu} \otimes x \mathbf{e}_2 \otimes y^{-1} \mathbf{f}_1 \otimes q^{\mathbf{h}_2 - \nu})$$

$$\times q^{\frac{(\mathbf{h}_1 + \mu) \otimes 1 \otimes (\mathbf{h}_1 + \nu) \otimes 1}{2} + \frac{1 \otimes (\mathbf{h}_2 - \mu) \otimes ((\mathbf{h}_1 + \nu) \otimes 1 + 1 \otimes (\mathbf{h}_2 - \nu))}{2}}, \quad (\text{A13})$$

where $\mathbf{e}\mathbf{v}_y^{(2)}$ is defined by (A2) replacing (μ, x) with (ν, y) .

Appendix B: q-analogue of H'Adamar formula

Here we mention formulas derived from the H'Adamar identity [18]

$$\exp_q(\mathbf{A}) \mathbf{B} \exp_q^{-1}(\mathbf{A}) = \sum_{k=0}^{\infty} \frac{\mathbf{B}_k}{(k)_q!}, \quad (\text{B1})$$

where $\mathbf{B}_0 = \mathbf{B}$, $\mathbf{B}_{k+1} = [\mathbf{A}, \mathbf{B}_k]_{q^k}$. Set $\mathbf{O} = \exp_{q^{-2}}(\lambda \mathbf{e}_1 \otimes \mathbf{f}_2)$. To simplify the notation, we will abbreviate $\mathbf{o}_1 \otimes 1$ and $1 \otimes \mathbf{o}_2$ as $\mathbf{o}_1$ and $\mathbf{o}_2$ respectively for any element $\mathbf{o}$ of the oscillator algebra. Then we obtain (as formal series on $\mathbf{e}_1 \mathbf{f}_2$)

$$\mathbf{O}^{-1} \mathbf{e}_1 \mathbf{O} = \mathbf{e}_1, \quad \mathbf{O}^{-1} \mathbf{e}_2 \mathbf{O} = \mathbf{e}_2 - (1 - q\lambda^2 \mathbf{e}_1 \mathbf{f}_2)^{-1} \mathbf{e}_1 q^{-\mathbf{h}_2},$$

$$\mathbf{O}^{-1} \mathbf{f}_1 \mathbf{O} = \mathbf{f}_1 - (1 - q\lambda^2 \mathbf{e}_1 \mathbf{f}_2)^{-1} \mathbf{f}_2 q^{\mathbf{h}_1}, \quad \mathbf{O}^{-1} \mathbf{f}_2 \mathbf{O} = \mathbf{f}_2, \quad (\text{B2})$$

$$\mathbf{O}^{-1} q^{\alpha \mathbf{h}_1} \mathbf{O} = \frac{(q^{2\alpha+1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^2)_\infty}{(q\lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^2)_\infty} q^{\alpha \mathbf{h}_1}, \quad \mathbf{O}^{-1} q^{\alpha \mathbf{h}_2} \mathbf{O} = \frac{(q^{-2\alpha+1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^2)_\infty}{(q\lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^2)_\infty} q^{\alpha \mathbf{h}_2},$$

where $\alpha \in \mathbb{C}$, $(a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k)$. From the last two identities, one can derive the following relations.

$$\mathbf{O}^{-1} q^{\alpha(\mathbf{h}_1 + \mathbf{h}_2)} \mathbf{O} = q^{\alpha(\mathbf{h}_1 + \mathbf{h}_2)}, \quad \mathbf{O}^{-1} q^{\pm \frac{\mathbf{h}_1 - \mathbf{h}_2}{2}} \mathbf{O} = (1 - q^{\pm 1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2)^{\mp 1} q^{\pm \frac{\mathbf{h}_1 - \mathbf{h}_2}{2}}. \quad (\text{B3})$$

One can calculate the inverse of (B2).

$$\begin{aligned} \mathbf{O} \mathbf{e}_1 \mathbf{O}^{-1} &= \mathbf{e}_1, & \mathbf{O} \mathbf{e}_2 \mathbf{O}^{-1} &= \mathbf{e}_2 + \mathbf{e}_1 q^{-\mathbf{h}_2}, \\ \mathbf{O} \mathbf{f}_1 \mathbf{O}^{-1} &= \mathbf{f}_1 + \mathbf{f}_2 q^{\mathbf{h}_1}, & \mathbf{O} \mathbf{f}_2 \mathbf{O}^{-1} &= \mathbf{f}_2, \\ \mathbf{O} q^{\alpha \mathbf{h}_1} \mathbf{O}^{-1} &= \frac{(q^{2\alpha-1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^{-2})_\infty}{(q^{-1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^{-2})_\infty} q^{\alpha \mathbf{h}_1}, & \mathbf{O} q^{\alpha \mathbf{h}_2} \mathbf{O}^{-1} &= \frac{(q^{-2\alpha-1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^{-2})_\infty}{(q^{-1} \lambda^2 \mathbf{e}_1 \mathbf{f}_2; q^{-2})_\infty} q^{\alpha \mathbf{h}_2}. \end{aligned} \quad (\text{B4})$$

Acknowledgments

The authors would like to thank Vladimir Bazhanov for useful discussions. The work of S.K. was supported by RFBR grant 14-01-00547. The work of Z.T. was supported by the Australian Research Council. A main part of this paper was written when he was at the department of theoretical physics, research school of physics and engineering, Australian national university.

References

- [1] R.J. Baxter, Partition function of the eight-vertex lattice model, *Ann. Phys.* 70 (1972) 193-228.
- [2] V.V. Bazhanov, S.L. Lukyanov, A.B. Zamolodchikov, Integrable Structure of Conformal Field Theory III. The Yang-Baxter Relation, *Commun.Math.Phys.* 200 (1999) 297-324 [arXiv:hep-th/9805008].
- [3] V.V. Bazhanov, A.N. Hibberd, S.M. Khoroshkin, Integrable structure of W_3 Conformal Field Theory, Quantum Boussinesq Theory and Boundary Affine Toda Theory, *Nucl. Phys. B* 622 (2002) 475–547 [arXiv:hep-th/0105177];
P. P. Kulish, A. M. Zeitlin, “Superconformal field theory and SUSY N=1 KDV hierarchy II: The Q-operator,” *Nucl. Phys. B* **709** (2005) 578 [hep-th/0501019];
H. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, Hidden Grassmann structure in the XXZ model, *Commun. Math. Phys.* 272 (2007) 263-281 [arXiv:hep-th/0606280];
T. Kojima, “The Baxter’s Q-operator for the W-algebra W_N ,” *J.Phys.A: Math. Theor.* 41 (2008) 355206 [arXiv:0803.3505 [nlin.SI]];
I. C.-H. Ip, A. M. Zeitlin, Q-operator and fusion relations for $C_q^{(2)}(2)$, arXiv:1312.4063 [math.QA].
- [4] H. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, Hidden Grassmann Structure in the XXZ Model II: Creation Operators, *Commun. Math. Phys.* 286 (2009) 875-932 [arXiv:0801.1176 [hep-th]].

- [5] V.V. Bazhanov, Z. Tsuboi: Baxter's Q-operators for supersymmetric spin chains, Nucl. Phys. B 805 [FS] (2008) 451-516 [arXiv:0805.4274 [hep-th]].
- [6] H. Boos, F. Göhmann, A. Klümper, K.S. Nirov, A.V. Razumov, Exercises with the universal R-matrix, J. Phys. A: Math. Theor. 43 (2010) 415208 [arXiv:1004.5342 [math-ph]]; On the universal R-matrix for the Izergin-Korepin model, J. Phys. A44 (2011) 355202 [arXiv:1104.5696 [math-ph]]; Universal R-matrix and functional relations, arXiv:1205.1631 [math-ph]; Quantum groups and functional relations for higher rank, arXiv:1312.2484 [math-ph].
- [7] Z. Tsuboi, Asymptotic representations and q-oscillator solutions of the graded Yang-Baxter equation related to Baxter Q-operators, arXiv:1205.1471 [math-ph].
- [8] D. Hernandez, M. Jimbo, Asymptotic representations and Drinfeld rational fractions, Compos. Math. 148 (2012) 1593-1623 [arXiv:1104.1891 [math.QA]]; E. Frenkel, D. Hernandez, Baxter's Relations and Spectra of Quantum Integrable Models, arXiv:1308.3444 [math.QA].
- [9] S. E. Derkachov, Factorization of the R-matrix.I, J. Math. Sci. 143 (2007) 2773-2790 [arXiv:math/0503396]; Factorization of the R-matrix.II, J. Math. Sci. 143 (2007) 2791-2805 [arXiv:math/0503410]; Factorization of R-matrix and Baxter's Q-operator, J. Math. Sci. 151 (2008) 2848-2858 [arXiv:math/0507252]; S. E. Derkachov, A. N. Manashov, Factorization of R-matrix and Baxter Q-operators for generic $sl(N)$ spin chains, J. Phys. A42 (2009) 075204 [arXiv:0809.2050 [nlin.SI]]; D. Chicherin, S. Derkachov, D. Karakhanyan, R. Kirschner, Baxter operators with deformed symmetry, Nucl. Phys. B [FS] 868 (2013) 652-683 [arXiv:1211.2965 [math-ph]]; V. Bazhanov, R. Frassek, T. Lukowski, C. Meneghelli, M. Staudacher, Baxter Q-Operators and Representations of Yangians, Nucl.Phys. B850 (2011) 148-174 [arXiv:1010.3699 [math-ph]]; R. Frassek, T. Lukowski, C. Meneghelli, M. Staudacher, Oscillator Construction of $su(n|m)$ Q-Operators, Nucl. Phys. B 850 (2011) 175-198 [arXiv:1012.6021 [math-ph]]; "Baxter Operators and Hamiltonians for 'nearly all' Integrable Closed $gl(n)$ Spin Chains," Nucl. Phys. B 874 [PM] (2013) 620-646 [arXiv:1112.3600 [math-ph]].
- [10] V. Bazhanov, T. Lukowski, C. Meneghelli, M. Staudacher, A Shortcut to the Q-Operator, J.Stat.Mech.1011:P11002,2010 [arXiv:1005.3261 [hep-th]].
- [11] V. Drinfeld, Hopf algebras and the quantum Yang-Baxter equations, Sov. Math. Dokl. 32 (1985) 254-258.
- [12] S. M. Khoroshkin and V. N. Tolstoy, The uniqueness theorem for the universal R-matrix, Lett. Math. Phys. 24 (1992) 231-244.
- [13] V. N. Tolstoy, S. M. Khoroshkin, The universal R-matrix for quantum untwisted affine Lie algebras, Funct. Anal. Appl. 26 (1992) 69-71;

- S. Khoroshkin, A. A. Stolin, V.N. Tolstoy, Gauss decomposition of trigonometric R-matrices, arXiv:hep-th/9404038; Generalized Gauss decomposition of trigonometric R-matrices, Mod.Phys.Lett. A10 (1995) 1375-1392.
- [14] M. Chaichian and P. Kulish, “Quantum Lie Superalgebras and q-Oscillators,” Phys. Lett. B **234**, 72-80 (1990).
 - [15] V.V. Bazhanov, S.M. Khoroshkin, (2001) unpulished.
 - [16] V.V. Bazhanov, Z. Tsuboi: talks at conferences in 2007, which include the following two: La 79eme Rencontre entre physiciens theoriciens et mathematiciens “Supersymmetry and Integrability”, IRMA Strasbourg, June, 2007; “Workshop and Summer School: From Statistical Mechanics to Conformal and Quantum Field Theory”, the university of Melbourne, January, 2007.
 - [17] Z. Tsuboi, Solutions of the T -system and Baxter equations for supersymmetric spin chains, Nucl. Phys. B 826 [PM] (2010) 399-455 [arXiv:0906.2039 [math-ph]].
 - [18] S. M. Khoroshkin and V. N. Tolstoy, Universal R-matrix for quantized (super)algebras, Commun. Math. Phys. 141 (1991) 599-617.